

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Karl Reinhard Papers/Publications

Natural Resources, School of

2018

The Paleoepidemiology of *Enterobius vermicularis* (Nemata: Oxyuridae) Among the Loma San Gabriel at La Cueva de los Muertos Chiquitos (600–800 CE), Rio Zape Valley, Durango, Mexico

Johnica J. Morrow

Karl Reinhard

Follow this and additional works at: <https://digitalcommons.unl.edu/natresreinhard>



Part of the [Archaeological Anthropology Commons](#), [Ecology and Evolutionary Biology Commons](#), [Environmental Public Health Commons](#), [Other Public Health Commons](#), and the [Parasitology Commons](#)

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Karl Reinhard Papers/Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

The Paleoepidemiology of *Enterobius vermicularis* (Nemata: Oxyuridae) Among the Loma San Gabriel at La Cueva de los Muertos Chiquitos (600–800 CE), Rio Zape Valley, Durango, Mexico

Authors: Morrow, Johnica J., and Reinhard, Karl J.

Source: Comparative Parasitology, 85(1) : 27-33

Published By: The Helminthological Society of Washington

URL: <https://doi.org/10.1654/1525-2647-85.1.27>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

The Paleoepidemiology of *Enterobius vermicularis* (Nemata: Oxyuridae) Among the Loma San Gabriel at La Cueva de los Muertos Chiquitos (600–800 CE), Rio Zape Valley, Durango, Mexico

JOHNICA J. MORROW^{1,2,3} AND KARL J. REINHARD¹

¹ Palynology and Pathoecology Laboratory, School of Natural Resources, University of Nebraska-Lincoln, 3310 Holdrege Street, Lincoln, Nebraska, 68583-0962, U.S.A. (email: kreinhard1@mac.com and

² Department of Physical and Life Sciences, Chadron State College, 1000 Main St., Chadron, Nebraska, 69337, U.S.A. (email: jmorrow@csc.edu)

ABSTRACT: One hundred coprolites excavated from La Cueva de los Muertos Chiquitos (600–800 CE) in the Rio Zape Valley of present-day Durango, Mexico, were examined for the presence of helminth eggs utilizing standard archaeoparasitological techniques. Eggs of the human pinworm (*Enterobius vermicularis*) were recovered from 34 of the 100 coprolites examined. Eggs of parasites were photographed and measured before egg concentration values were calculated for each positive sample. Egg concentration values demonstrated an overdispersed pattern of distribution among the samples (66% uninfected, 25% less than 100 eggs/g, 8% between 100 and 500 eggs/g, and 1% more than 500 eggs/g). Given that only 5% of infected hosts in modern cases of human enterobiasis pass the eggs of parasites in their stools, the recovery of *E. vermicularis* eggs in 34% of the coprolites supports the conclusion that virtually all of the individuals utilizing the site during the coprolite depositional time frame likely were infected with this parasite. These data are discussed in light of other studies of prehistoric human enterobiasis.

KEY WORDS: archaeoparasitology, coprolite, Durango, Mexico, *Enterobius vermicularis*, La Cueva de los Muertos Chiquitos, paleoepidemiology, pathoecology, pinworm.

The pinworm, *Enterobius vermicularis*, is a common human parasite with evidence showing a host-parasite association going back as far as 8,000 years (Hugot et al., 1999). In a modern context, infection with this parasite occurs commonly among humans with prevalences among children, childcare workers, and institutionalized persons being reported as high as 50% (Burkhart and Burkhart, 2005). The oldest human pinworm eggs recovered from human remains in North America were reported from Danger Cave, an archaeological site in Utah that dates to about 10,200 calendar years ago (Rhode and Madsen, 1998; Goebel et al., 2007). Eggs identified as *E. vermicularis* have been recovered from many other North American sites that predate European contact of the thirteenth century AD, which suggests that this parasite was established in the Americas alongside their human hosts following migrations into the continent (Reinhard et al., 2016).

Evidence indicates that in parts of the prehistoric Americas, *E. vermicularis* was seemingly epidemic. Bearing in mind that only 5% of infected individuals release eggs in their stools, it is remarkable that pinworm eggs are often recovered from co-

prolites excavated from sites in the Greater Southwest in numbers exceeding 10% (Fry, 1974; Faulkner et al., 1989; Reinhard, 1992; Fugassa et al., 2011; Jiménez et al., 2012; Reinhard et al., 2016). Subsistence strategies and dwelling types have quantifiable effects on pinworm prevalences among ancient peoples (Hugot et al., 1999), and agricultural populations were more frequently found to have pinworms than hunter-gatherers, largely due to the impacts of sedentism (Reinhard, 1988). Ancient dwellings, many of which were built within caves, served as nidi for these parasites (Reinhard and Bryant, 2008), while the living habits of agriculturalists' were conducive to shifts in human parasitism in terms of both diversity and intensity of infection, which is reflected in archaeoparasitological analyses of coprolites (Reinhard, 1988).

Archaeoparasitological data can be utilized to examine the paleoepidemiology of ancient parasitism in ways similar to modern epidemiological studies (Han et al., 2003; Seo et al., 2008, 2014). Early paleoepidemiological studies of parasites began with Fry's examination of parasite prevalence in the Great Basin spanning nearly 10,000 years (Fry, 1977). Further examinations of the changes in parasitism related to Colorado Plateau hunter-gatherer/agriculturalist transitions were studied by Reinhard beginning a decade

³ Corresponding author.

later (Reinhard, 1988, 1992), and paleoepidemiological studies in North American resumed only recently (Fugassa et al., 2011; Jiménez et al., 2012; Morrow and Reinhard, 2016).

In the present study, 100 coprolites excavated from La Cueva de los Muertos Chiquitos (CMC) were examined for the presence of *E. vermicularis* eggs. CMC is a cave located within the Rio Zape Valley, a region approximately 18 kilometers southeast of Guanaceví in Durango, Mexico (Jiménez et al., 2012; Morrow and Reinhard, 2016). This valley houses a series of caves once utilized by people of the Loma San Gabriel culture between 1,200 and 1,400 years ago (Brooks et al., 1962; Foster, 1986). This area represents an archaeological transition zone between the northernmost edge of Mesoamerica and the greater American Southwest (Kelley, 1956, 1971; Brooks and Brooks, 1980). Excavations of CMC recovered 21 skeletons of children, ranging between several months and three years of age, and more than 500 well-preserved coprolites (Brooks et al., 1962; Brooks and Brooks, 1978, 1980). These coprolites were sealed beneath adobe floors, which inhibited the influence of certain taphonomic agents, such as water, wind, flies, and scavengers (Morrow and Reinhard, 2016; Morrow et al., 2016).

Parasitism among the Loma San Gabriel has been predominately studied through the analysis of CMC coprolites (Jiménez et al., 2012; Cleeland et al., 2013; Morrow and Reinhard, 2016), and previous study of 36 CMC coprolites reported a 44% prevalence of *E. vermicularis* eggs (Jiménez et al., 2012). Herein, the recovery of *E. vermicularis* eggs from examinations of an additional 100 CMC coprolites are reported and the paleoepidemiology of this parasite at CMC is discussed.

MATERIALS AND METHODS

Coprolites from La Cueva de los Muertos Chiquitos (CMC) archived within the Pathoecology Laboratory in the School of Natural Resources at the University of Nebraska were used for the present study. This is a filtered-air, positive-pressure facility that is free of contaminants smaller than a micrometer. A total of 100 coprolites from this collection were given an individual analysis identification number, weighed, photographed, and placed into individually labeled plastic zippered bags. Many of these coprolites had been stored in the same bags as other coprolites, which gave these artifacts the potential to contaminate one another

with the mechanical transfer of eggs, especially because these materials have been transported repeatedly since their excavation in the early 1960s. To reduce the retention of potentially contaminate eggs, the surfaces of each of the coprolites were abraded using cleaned, stiff brushes. To prevent modern contamination of material, nitrile gloves were worn during cleaning. Dusts from the abraded surfaces of these coprolites were retained in their primary analysis bags, and each coprolite was placed into a clean secondary analysis bag.

A subsample was extracted from each of the coprolites and weighed for standard analysis of parasites. These subsamples were rehydrated with 0.5% trisodium phosphate for 24 hr prior to disaggregation utilizing a magnetic stirring apparatus. The color of the rehydrated material was recorded. Samples were subsequently treated with 1 tablet/g of sample containing the spores of *Lycopodium* (Batch no. 124961, Department of Quaternary Geology, University of Lund, Lund, Sweden; containing approximately 12,500 spores/tablet). These tablets were dissolved using hydrochloric acid (HCl) in sterilized 50 ml plastic beakers and rinsed into each beaker of disaggregated material (Stockmarr, 1971).

The method for using *Lycopodium* to quantify the eggs of parasites from archaeological materials was first developed by Warnock and Reinhard (1992) and has become a standard quantification technique among archaeo- and paleoparasitologists (Martinson et al., 2003; Reinhard et al., 2008, 2012; Reinhard and Urban, 2003; Santoro et al., 2003; Sianto et al., 2005; Kumm et al., 2010; Fugassa et al., 2011; Jiménez et al., 2012; Searcey et al., 2013). By spiking samples with a known amount of *Lycopodium* spores, subsequent microfossil (e.g., eggs, pollen grains, and phytoliths) concentrations can be determined. During light microscopy analysis of the samples, the number of *Lycopodium* spores recovered is counted along with the numbers of microfossils observed. A microfossil concentration formula is then used to estimate the number of microfossils per gram of material as follows: Microfossil concentration = $[(p/m) \times a]/w$, where p is the number of microfossils counted, m is the number of marker grains (*Lycopodium* spores) counted, a is the number of *Lycopodium* spores added to the sample, and w is the total weight of the subsample prior to rehydration.

After dosing with dissolved *Lycopodium* spore tablets, samples were passed through a 250 μ m mesh screen to separate macroscopic remains (material

larger than 250 μm) from microscopic remains (material smaller than 250 μm). Macroscopic remains were transferred to clean, appropriately labeled filter paper and allowed to dry.

The microscopic remains obtained following screening were concentrated into 50 ml plastic, screw-cap, graduated centrifuge tubes via repeated centrifugation. Examination slides of each sample were created by mixing a small amount of concentrated material with a drop of glycerin onto a clean, glass microscope slide and topping with a 22 mm \times 22 mm cover slip. Slides were examined using a Nikon compound microscope. Any eggs encountered were photographed using a Sony Cybershot 18.2 megapixel camera and measured using an integrated ocular micrometer. Each egg was photographed and measured except in cases where eggs were damaged or folded and accurate measurements could not be obtained (Fig. 2). Subsequently, parasite egg concentration values were calculated for each of the 34 samples containing the eggs of *E. vermicularis*. All measurements are in microns unless otherwise noted.

The unprocessed half of each sample was saved in its designated secondary analysis bag and returned to the CMC collection of material in the Pathoecology Laboratory. Additionally, half of each processed subsample was archived and labeled as a preparation of parasite eggs using standard archaeoparasitological archival techniques. The other halves of the processed subsamples were preserved in 95% EtOH for future processing for palynological analysis.

RESULTS

Many of the eggs observed contained developed larvae, and microscopic analysis revealed evidence of *E. vermicularis* in 34/100 samples. The average egg concentration value for coprolites containing eggs was 157 eggs/g. Egg concentration values ranged from 36 eggs/g to 1,901 eggs/g. Concentration values expressed in eggs/coprolite rather than eggs/g ranged from 372 eggs/coprolite to 2,985 eggs/coprolite, with an average of 373 eggs/coprolite. Recovered eggs ranged in length from 49.0 to 63.7 and in width from 19.6 to 31.85. On average, the eggs recovered from CMC coprolites measured 56 \times 29 (Morrow, 2016, unpublished dissertation, University of Nebraska–Lincoln, Lincoln, Nebraska, U.S.A.).

When accounting for all of the coprolites examined in the present study, a pattern of overdispersion with regard to egg distribution emerges (Figs. 1, 2). No eggs

were recovered from 66% of the coprolites. Of the 34% that did contain the eggs of this parasite, the majority of coprolites presented with low concentrations of eggs, while only a few presented with high concentrations of eggs (Figs. 1, 2).

DISCUSSION

The results of the present archaeoparasitological analysis show that *E. vermicularis* was present among the Loma San Gabriel. Although a few of the eggs recovered were fractured, most of the eggs found among these samples were relatively well preserved. Photographs and measurements of recovered eggs were used to positively identify these eggs as *E. vermicularis*. The majority of the eggs fell within known parameters for *E. vermicularis*, which are 50–60 in length by 20–32 in width (Thienpoint et al., 1979), and those eggs that we observed that fell outside these parameters were only slightly outside of this range (i.e., 49 length or 19.6 width). These slightly smaller eggs could be the result of taphonomic changes as opposed to being authentically smaller in size (Morrow et al., 2016).

Given that a single gravid female *E. vermicularis* deposits approximately 4,600–16,000 eggs/day (Kliks, 1990; Ferreira et al., 1997; Roberts et al., 2012), these data indicate that the depositors of *E. vermicularis* positive coprolites were only lightly infected with these parasites (mean = 157 eggs/g; range = 54–2,985 eggs/coprolite). However, the eggs of this parasite are light weight and easily aerosoled, and may adhere to surfaces within a residence. These eggs may be transmitted via inhalation or ingestion in this manner (Hugot et al., 1999), and individuals may also become autoinfected. Though eggs are deposited within the intestine (Garcia, 2009), they are not laid within the feces themselves (Caldwell, 1982). For this reason, only 5–15% of modern cases of enterobiasis report the recovery of eggs from fecal examinations (Cook, 1994; Brooker and Bundy, 2009). Thus, despite the low concentrations of eggs recovered in the present study as compared to the fecundity of modern pinworms, the recovery of these eggs from 34% of CMC coprolites suggests that much of the population utilizing the site likely was infected. This further tells us that CMC was either occupied regularly enough to sustain the transmission cycle among different groups of Loma San Gabriel using the site, or that those who used the site perpetuated the life cycle among one another at other, more permanent sites of residence. Parasites like *E. vermicularis* proliferate in human populations with




Figure 1. Egg concentrations from La Cueva de los Muertos Chiquitos coprolites infected with *Enterobius vermicularis*.

somewhat sedentary life styles. This is a parasite often associated with overcrowding that spreads easily within continuously populated domiciles. Though it is not known exactly how many individuals are represented by the 100 coprolites examined, an idea of the population profile was recovered from dental impressions of quids by Hammerl et al. (2015). Quids are expectorated masses of fibers that are common in archaeological sites used by people dependent on high-fiber plant resources. Analysis of 50 quids revealed dental

impressions showing that the site was used by all age groups (Hammerl et al., 2015), with 49 distinct dental arcades from 49 people evident from the dental casts. These dental data suggest that a relatively large group of people, consistent with a permanent village, used the site. Additionally, 21 children younger than three years of age were buried at the site, which suggests that a larger population used the site. Given these data, it is estimated that the site was used by 100–250 people. In short, the life cycles of these parasites were being per-




Figure 2. The concentrations and distribution of *Enterobius vermicularis* eggs among La Cueva de los Muertos Chiquitos coprolites.

petuated in those utilizing CMC approximately 1,300 years ago.

The overdispersed nature of the data presented herein is commonly encountered in modern parasitological systems (Crofton, 1971; Croll and Ghadirian, 1981; Crompton et al., 1984; Anderson and May, 1985; Shaw et al., 1998). While overdispersion is commonly discussed in reference to modern parasitism, it has rarely been discussed from an archaeoparasitological perspective. At CMC, the majority (66%) of coprolites did not contain *E. vermicularis* eggs, while the majority of the 34% of coprolites that did contain these eggs presented with very low egg concentrations, and only a small percentage of coprolites presented with high egg concentrations (Figs. 1, 2). Because it is not known precisely how many human individuals are represented by the 100 coprolites examined, it is not possible to definitively state that a negative binomial pattern of distribution of these eggs existed within this host population. Similarly, because many infected individuals do not pass the eggs of *E. vermicularis* in their stool, it is difficult to assess how many individuals were actively infected with this parasite. Future analyses of the remaining material from these coprolites utilizing aDNA extraction methods could shed light on the nature of *E. vermicularis* distribution among the Loma San Gabriel at La Cueva de los Muertos Chiquitos. Such future analyses should seek to assess the number of individual hosts represented by comparing human aDNA among the samples and should examine the coprolites for molecular evidence of *E. vermicularis* to more accurately assess the rates of infectivity among this population.

Antelope House is an archaeological site occupied around 900 years ago and was a permanent village constructed in a large rockshelter. Previous analysis of Antelope House yielded a 24% prevalence of *E. vermicularis* eggs among 180 coprolites (Hugot et al., 1999), a number similar to what we observed at CMC (20% prevalence of *E. vermicularis* eggs among 100 coprolites). Overall, Hugot et al. (1999) examined coprolites from 10 archaeological sites, dated from 8000 BC to 1300 AD, for the presence of *E. vermicularis* eggs. Their results indicate that only 1.8% of hunter-gatherer sites in caves or rock shelters were positive for pinworm eggs, and no pinworm eggs were recovered from cave sites without stone-walled villages, while cave sites with stone-walled villages presented with a prevalence of 19% among examined coprolites.

The results of the present study are similar to those reported by Jiménez et al. (2012), who examined 36

coprolites from CMC and found *E. vermicularis* prevalence to be 26% utilizing a flotation technique and 35% utilizing a sedimentation technique. By increasing the sample size to 100 CMC coprolites and using a standard rehydration, disaggregation, and screening technique, we report a 34% prevalence of *E. vermicularis* eggs among CMC coprolites. Combining the results of these studies, *E. vermicularis* eggs have now been reported from 50 of 136 CMC coprolites (36.8% prevalence among CMC coprolites).

The data presented herein expand our understanding of prehistoric human enterobiasis in the New World and are especially significant for establishing the paleoepidemiological patterns of pinworm infections among a transitional culture such as the Loma San Gabriel. Future analyses of both host and parasite aDNA extracted from CMC coprolites could further elucidate paleoepidemiological patterns.

ACKNOWLEDGMENTS

The authors would like to thank the School of Natural Resources, University of Nebraska–Lincoln, for the purchase of the research microscope used during these analyses.

LITERATURE CITED

- Anderson, R., and R. May. 1985. Helminth infections of humans: mathematical models, population dynamics, and control. *Advances in Parasitology* 24:1–101.
- Brooker, S., and D. A. P. Bundy. 2009. Soil-transmitted Helminths (Geohelminths). Pages 1515–1549 in G. C. Cook and A. I. Zumla, eds. *Manson's Tropical Diseases*. 22nd Ed., Saunders Elsevier.
- Brooks, R. H., L. Kaplan, H. C. Cutler, and T. W. Whitaker. 1962. Plant material from a cave on the Rio Zape, Durango, Mexico. *American Antiquity* 27:356–369.
- Brooks, S. T., and R. H. Brooks. 1978. Paleoepidemiology as a possible interpretation of multiple child burials near Zape Chico, Durango, Mexico. Pages 96–101 in J. C. Kelley, C. L. Riley, and B. C. Hedrick, eds. *Across the Chichimec Sea: Papers in Honor of J. Charles Kelley*. Southern Illinois University Press, Carbondale, Illinois.
- Brooks, S. T., and R. H. Brooks. 1980. Cranial deformation: Possible evidence of Pochteca trading movements. *Transactions of the Illinois State Academy of Sciences* 72:1–12.
- Burkhart, C. N., and C. G. Burkhart. 2005. Assessment of frequency, transmission, and genitourinary complications of enterobiasis (pinworms). *International Journal of Dermatology* 44(10):837–840.
- Caldwell, J. P. 1982. Pinworms (*Enterobius vermicularis*). *Canadian Family Physician* 28:306–309.
- Cleeland, L. M., M. V. Reichard, R. Y. Tito, K. J. Reinhard, and C. M. Lewis, Jr. 2013. Clarifying prehistoric parasitism from a contemporary morphological and

- molecular approach. *Journal of Archaeological Science* 40:3060–3066.
- Cook, G. C.** 1994. *Enterobius vermicularis* infection. *Gut* 35(9):1159–62.
- Crofton, H. D.** 1971. A quantitative approach to parasitism. *Parasitology* 62:179–193.
- Croll, N., and E. Ghadirian.** 1981. Wormy persons: contributions to the nature and patterns of overdispersion with *Ascaris lumbricoides*, *Ancylostoma duodenale*, *Necator americanus* and *Trichuris trichiura*. *Tropical and Geographical Medicine* 33:241–248.
- Crompton, D., A. Keymer, and S. Arnold.** 1984. Investigating over-dispersion: *Moniliformis* (Acanthocephala) and rats. *Parasitology* 88:317–331.
- Faulkner, C. T., S. Patton, and S. S. Johnson.** 1989. Prehistoric parasitism in Tennessee: evidence from the analysis of desiccated fecal material collected from Big Bone Cave, Van Buren County, Tennessee. *Journal of Parasitology* 75:461–463.
- Ferreira, L. F., K. J. Reinhard, A. Araújo, and L. Camillo-Coura.** 1997. Paleoparasitology of oxyuriasis. *Anales de la Real Academia Nacional de Medicina* 157:20–24.
- Foster, M. S.** 1986. The Weicker site: A Loma San Gabriel hamlet in Durango, Mexico. *Journal of Field Archaeology* 13:7–19.
- Fry, G. F.** 1974. Ovum and parasite examination of Salts Cave human paleofeces. Page 61 in P. J. Watson, ed. *Archeology of the Mammoth Cave Area*. Academic Press, New York.
- Fry, G. F.** 1977. Analysis of prehistoric coprolites from Utah. *University of Utah Anthropological Papers*, Vol. 97. University of Utah Press, Salt Lake City, Utah.
- Fugassa, M. H., K. J. Reinhard, K. L. Johnson, M. Vieira, and A. Araújo.** 2011. Parasitism of prehistoric humans and companion animals from Antelope Cave, Mojave County, Northwest Arizona. *Journal of Parasitology* 97(5):862–867.
- García, L. S.** 2009. Practical guide to diagnostic parasitology. *American Society for Microbiology* 246–247.
- Goebel, T., K. Graf, and D. Rhode.** 2007. The Paleoindian occupations at Bonneville Estates Rockshelter, Danger Cave, and Smith Creek Cave (eastern Great Basin, USA): interpreting their radiocarbon chronologies. *Archaeopress*, Pages 147–161 in M. Kornfeld, S. Vasilev, and L. Motti, eds. *On Shelter's Ledge: Histories, Theories and Methods of Rockshelter Research*. BAR International Series 1655. Oxford, UK.
- Hammerl, E. E., M. A. Baier, and K. J. Reinhard.** 2015. Agave chewing and dental wear: evidence from quids. *PLoS One* 10(7):e0133710.
- Han, E.-T., S.-M. Guk, J.-L. Kim, H.-J. Jeong, S.-N. Kim, and J.-Y. Chai.** 2003. Detection of parasite eggs from archaeological excavations in the Republic of Korea. *Memórias do Instituto do Oswaldo Cruz* 98:123–126.
- Hugot, J. P., K. J. Reinhard, S. L. Gardner, and S. Morand.** 1999. Human enterobiasis in evolution: origin, specificity and transmission. *Parasite* 6(3): 201–208.
- Jiménez, F. A., S. L. Gardner, A. Araújo, M. H. Fugassa, R. H. Brooks, E. Racz, and K. J. Reinhard.** 2012. Zoonotic and human parasites of inhabitants of Cueva de Los Muertos Chiquitos, Rio Zape Valley, Durango, Mexico. *Journal of Parasitology* 98(2): 304–309.
- Kelley, J. C.** 1956. Settlement patterns in North-Central Mexico. Pages 128–129 in G. R. Willey, ed. *Prehistoric Settlement Patterns in the New World*. Viking Fund Publications in Anthropology, New York.
- Kelley, J. C.** 1971. Archaeology of the northern frontier: Zatecas and Durango. Page 768–801 in G. F. Ekholm and I. Bernal, eds. *Handbook of Middle American Indians*, Vol. 11. *Archaeology of Northern Mesoamerica*, Part 2. University of Texas Press, Austin, Texas.
- Kliks, M. M.** 1990. Helminths as heirlooms and souvenirs: a review of New World paleoparasitology. *Parasitology Today* 6:93–100.
- Kumm, K., K. J. Reinhard, A. Araújo, and D. Piombino-Mascali.** 2010. Archaeoparasitological investigation of a mummy from Sicily (18th–19th century AD). *Anthropologie* 48:177–184.
- Martinson, E., K. J. Reinhard, J. E. Buikstra, and K. Dittmar.** 2003. Pathoecology of Chiribaya parasitism. *Memorias do Instituto do Oswaldo Cruz* 98:195–205.
- Morrow, J. J., J. Newby, D. Piombino-Mascali, and K. J. Reinhard.** 2016. Taphonomic considerations for the analysis of parasites in archaeological materials. *International Journal of Paleopathology* 13:56–64.
- Morrow, J. J., and K. J. Reinhard.** 2016. *Cryptosporidium parvum* among Coprolites from La Cueva de los Muertos Chiquitos (600–800 AD), Rio Zape Valley, Durango, Mexico. *Journal of Parasitology* 102(4):429–435.
- Reinhard, K.** 1988. Cultural ecology of prehistoric parasitism on the Colorado Plateau as evidenced by coprology. *American Journal of Physical Anthropology* 77:355–366.
- Reinhard, K.** 1992. Patterns of diet, parasitism and anemia in prehistoric west North America. Pages 219–258 in P. Stuart-Macadam, and S. Kent, eds. *Diet, Demography and Disease: Changing Perspectives on Anemia*. Aldine de Gruyter, New York.
- Reinhard, K. J., and V. M. Bryant.** 2008. Pathoecology and the future of coprolite studies. Pages 199–216 in A. W. M. Stodder, ed. *Reanalysis and Reinterpretation in Southwestern Bioarchaeology*. Arizona State University Press, Tempe, Arizona.
- Reinhard, K., and Urban, O.** 2003. Diagnosing ancient diphyllbothriasis from Chinchorro mummies. *Memorias do Instituto do Oswaldo Cruz* 98:191–193.
- Reinhard, K. J., A. Araújo, and J. J. Morrow.** 2016. Temporal and spatial distribution of *Enterobius vermicularis* (Nemata:Oxyuridae) in the prehistoric Americas. *Korean Journal of Parasitology* 54(5):1–13.
- Reinhard, K. J., A. Araújo, L. Sianto, J. G. Costello, and K. Swope.** 2008. Chinese liver flukes in latrine sediments from Wong Nim's property, San Bernardino, California: archaeoparasitology of the Caltrans District Headquarters. *Journal of Parasitology* 94: 300–303.
- Reinhard, K. J., K. L. Johnson, S. LeRoy-Toren, K. Wieseman, I. Teixeira-Santos, and M. Vieira.** 2012. Understanding the pathoecological relationship between ancient diet and modern diabetes through coprolite analysis: a case example from Antelope Cave, Mojave County, Arizona. *Current Anthropology* 53: 506–512.
- Rhode, D., and D. B. Madsen.** 1998. Pine nut use in the early holocene and beyond: the Danger Cave archaeobotanical record. *Journal of Archaeological Science* 25: 1199–1210.

- Roberts, L., J. Janovy Jr., and S. Nadler.** 2012. *Foundations of Parasitology*, 9th ed. McGraw-Hill, New York, New York.
- Santoro, C., S. D. Vinton, and K. Reinhard.** 2003. Inca expansion and parasitism in the Lluta Valley: preliminary data. *Memorias do Instituto do Oswaldo Cruz* 98: 161–163.
- Searcey, N., K. J. Reinhard, S. L. Gardner, E. Egarter-Vigl, F. Maixner, D. Piombino-Mascali, A. Zink, W. Van Der Sanden, and R. Bianucci.** 2013. Parasitism of the Zweeloo Woman: microcoeliasis evidenced in a Roman Period bog mummy. *International Journal of Paleopathology* 3:224–228.
- Seo, M., C. S. Oh, J.Y. Chai, M. S. Jeong, S. W. Hong, Y.-M. Seo, and D. H. Shin.** 2014. The changing pattern of parasitic infection among Korean populations by paleoparasitological study of Joseon Dynasty mummies. *Journal of Parasitology* 100: 147–150.
- Seo, M., D. H. Shin, S.-M. Guk, C.S. Oh, E.-J. Lee, M. H. Shin, M. J. Kim, S. D. Lee, Y.-S. Kim, Y.S. Yi, M. Spigelman, and J.-Y. Chai.** 2008. *Gymnophalloides seoi* eggs from the stool of a 17th century female mummy found in Hadong, Republic of Korea. *Journal of Parasitology* 94:467–472.
- Shaw, D., B. Grenfell, and A. Dobson.** 1998. Patterns of macroparasite aggregation in wildlife host populations. *Parasitology* 117:597–610.
- Sianto, L., K. J. Reinhard, M. L. C. Gonçalves, and A. Araújo.** 2005. The finding of *Echinostoma* (Trematoda: Digenea) and hookworm eggs in coprolites collected from a Brazilian mummified body dated 600–1,200 years before present. *Journal of Parasitology* 91:972–975.
- Stockmarr, J.** 1971. Tablets with spores used in absolute pollen analysis. *Pollen et Spores* 13:615–621.
- Thienpont, D., F. Rochette, and O. F. J. Vanparijs.** 1979. Diagnosing helminthiasis through coprological examination. Janssen Research Foundation, Beerse, Belgium. 187 pp.
- Warnock, P. J., and K. J. Reinhard.** 1992. Methods for extracting pollen and parasite eggs from latrine soils. *Journal of Archaeological Science* 19: 261–264.